


# Physics of Hypernuclei as seen by an experimenter



# Outline of the lecture


- **interaction with strangeness**
  - the hyperon-nucleon interaction
- **hypernuclear formation in electroproduction**
  - the KAOS spectrometer at the accelerator MAMI-C
- **hypernuclear spectroscopy with stable heavy ion beams and rare isotope beams**
  - the HypHI experiment at GSI/FAIR
- **hypernuclear physics with anti-protons**
  - the Panda experiment at FAIR
- **summary of the activities**

# The hyperon-nucleon interaction

- **Y-N and Y-Y scattering:**
  - no hyperon target available:  
 $\tau$  (hyperon)  $\sim 10^{-10}$  s
  - very high energy hyperon beams:  
CERN WA89 and SELEX
  - low energy hyperons on nuclei:  
very poor hyperon beam profile

→ impossible to deduce precise Y-N data  
→ impossible to deduce any Y-Y data

- **hypernuclei:** nuclei with strange quarks
  - nuclear bound system with hyperon (Y)
  - a “laboratory” to study baryon-baryon interactions with strange quarks




[Fujiwara et al., nucl-th/0607013 ]

number of baryons  
 $N+Z+Y$

number of hyperons

Example  
 ${}^7_{\Lambda}\text{Li}$  ( ${}^6\text{Li}$ )




increasing strangeness

- only very few

stability

# Single $\Lambda$ -hypernuclei

$\Lambda$  Hypernuclear Chart  
(2005)



[Updated from: O. Hashimoto and H. Tamura, Prog. Part. Nucl. Phys. 57 (2006) 564]

# Single $\Lambda$ -hypernuclei

$$V_{\Lambda N}^{eff} = V_0 + \Delta(\vec{s}_\Lambda \cdot \vec{s}_N) + S_N(\vec{l}_{\Lambda N} \cdot \vec{s}_N) + S_\Lambda(\vec{l}_{\Lambda N} \cdot \vec{s}_\Lambda) + T(s_{12})$$

${}^7_\Lambda\text{Li} (3/2^+, 1/2^+)$

${}^7_\Lambda\text{Li} (5/2^+, 1/2^+)$

${}^9_\Lambda\text{Be} (3/2^+, 5/2^+)$

${}^{16}_\Lambda\text{O} (1^-, 0^-)$


$\Delta = 0.4 \text{ MeV}$

$S_N = -0.4 \text{ MeV}$

$S_\Lambda = -0.01 \text{ MeV}$


$T = 0.03 \text{ MeV}$

${}^7\text{Li} (\pi^+, K^+\gamma)$  KEK E419




${}^7_\Lambda\text{Li}$  PRL 84 (2000) 5963  
PRL 86 (2001) 1982  
PLB 579 (2004) 258

${}^9\text{Be} (K^-, \pi^-\gamma)$  BNL E930-1




PRL 88 (2002) 082501

${}^{16}\text{O} (K^-, \pi^-\gamma)$  BNL E930-2



PRL 93 (2004) 232501



# Impurity effects



VOLUME 86, NUMBER 10

PHYSICAL REVIEW LETTERS


5 MARCH 2001

## Measurement of the $B(E2)$ of ${}^7_{\Lambda}\text{Li}$ and Shrinkage of the Hypernuclear Size

K. Tanida,<sup>1</sup> H. Tamura,<sup>2</sup> D. Abe,<sup>2</sup> H. Akikawa,<sup>3</sup> K. Araki,<sup>2</sup> H. Bhang,<sup>4</sup> T. Endo,<sup>2</sup> Y. Fujii,<sup>2</sup> T. Fukuda,<sup>5</sup> O. Hashimoto,<sup>2</sup> K. Imai,<sup>3</sup> H. Hotchi,<sup>1</sup> Y. Kakiguchi,<sup>5</sup> J.H. Kim,<sup>4</sup> Y.D. Kim,<sup>6</sup> T. Miyoshi,<sup>2</sup> T. Murakami,<sup>3</sup> T. Nagae,<sup>5</sup> H. Noumi,<sup>5</sup> H. Outa,<sup>5</sup> K. Ozawa,<sup>2</sup> T. Saito,<sup>7</sup> J. Sasao,<sup>2</sup> Y. Sato,<sup>2</sup> S. Satoh,<sup>2</sup> R. I. Sawafta,<sup>8</sup> M. Sekimoto,<sup>5</sup> T. Takahashi,<sup>2</sup> L. Tang,<sup>9</sup>  
H. H



# Hyperon-hyperon interaction





$$B_{\Lambda\Lambda} / A \approx \dots + \dots$$

$$\Delta \dots = \dots - \dots$$

- 1963: Danysz *et al.*  $^{10}_{\Lambda\Lambda}\text{Be}$
- 1966: Prowse  $^6_{\Lambda\Lambda}\text{He}$
- 1991: KEK-E176  $^{13}_{\Lambda\Lambda}\text{B}$
- 2001: KEK-E373  $^6_{\Lambda\Lambda}\text{He}$
- 2001: AGS-E906  $^4_{\Lambda\Lambda}\text{H} ? (\sim 15)$

# Spectroscopy of $\Lambda\Lambda$ -hypernuclei

[E. Hiyama, M. Kamimura, T. Motoba, T. Yamada and Y. Yamamoto, Phys. Rev. 66 (2002), 024007]





[4-body cluster model for light nuclei]

many excited, particle stable states in double hypernuclei predicted  
 $\gamma$ -spectroscopy of these states is mandatory to study them

# $\Lambda\Lambda$ -Nuclei as Laboratory for H


H-Particle

[R.L. Jaffe (1977)]




T. Sakai, K. Shimizu, K. Yazaki  
 Prog.Theor.Phys.Suppl. 137 (2000) 121-145

# International hypernuclear network





# Strangeness reactions

exchange of strangeness



production of open strangeness



# Electroproduction




- electroproduction of hypernuclei
- neutron-richer single  $\Lambda$  hypernuclei
  - $\Lambda$  wave-function inside hypernucleus
  - large momentum transfer components

# Kinematic differences to meson induced reactions


- typical momentum transfers:  $\approx 300 - 600 \text{ MeV}/c$
- minimum momentum transfer for  $\theta_K = 0^\circ$
- energy and momentum transfer independent:

$$Q^2 = -q_\mu q^\mu = \omega^2 - \vec{q}^2$$

- momentum transfer  $\rightarrow 0$  for “magic momentum”
- minimum momentum transfer for  $\theta_\pi = 0^\circ$
- momentum distributions cannot be measured



[strangeness electroproduction ( $e, e' K^+$ )]




[strangeness exchange ( $K^-, \pi^-$ )]

hypernuclei: spectrometry of mesons at forward angles


# Extracting hypernuclear structure information

- cross sections calculated with harmonic oscillator potential and DWIA
- typical  $K^+$  angular distributions peaked at  $0^\circ$ , falling rapidly:




$\Delta E \sim 650$  keV; Core excited states  
Theoretical interpretation and publication in progress

[J. Reinhold (FIU), DNP Town Meeting, Dec. 2007]




[M. Sotona and S. Furullani, Prog. Theor. Phys. Suppl. 117, 151 (1994)]




»machines machines  
machines machines«  
»machines machines  
machines machines«  
  
»It's a machines world«

Queen, April 1984

# The »three« spectrometer facility at MAMI




# Transport of KAOS to Mainz in June 2003



compact, open yoke and extended pole face  
⇒ use as double spectrometer

# Two-arm spectrometer operation of KAOS




Physics of Hypernuclei as seen by an experimenter






# Kinematical optimisation using a *Figure Of Merit* for formation rate

$$\text{FOM} = S_{\Lambda} \times \Gamma \quad \text{with} \quad \Gamma = \frac{\alpha}{2\pi^2} \frac{E'}{E} \frac{k_{\gamma}}{Q^2} \frac{1}{1 - \epsilon}$$


[ $Q^2 = 0.01 \text{ GeV}^2/c^2$ ,  $W = 11.995 \text{ GeV}$ ,  $E = 1.50 \text{ GeV}$ ,  $E' = 0.650 \text{ GeV}$ ,  $\theta_e = 5.8^\circ$ ,  $p_K = 0.446 \text{ GeV}/c$ ,  $p_Y = 0.423 \text{ GeV}/c$ , and  $\theta_K = 5.5^\circ$ ]



# The detector packages for KAOS



Hadron arm completed in Feb 2008




## Timeline

- 2008: data taking for kaon production is running
- 2009: completion of KAOS as double spectrometer at  $0^\circ$
- 2009/10: data taking for hypernuclei production


# Challenges and prospects

special features of electro-production at MAMI-C (and JeffersonLab)


- better resolution compared to ( $\pi^+, K^+$ ) or ( $K^-, \pi^-$ )
- access to new isotopes of hypernuclei (converting p into  $\Lambda$ )
- measurements at different kaon angles map out different parts of the  $\Lambda$  momentum distribution
- unique with KAOS: double spectroscopy in a single spectrometer




expected mass resolution:  
 $\sigma_m = 275 \text{ keV}$




# Networking Activity SPHERE within the EU FP7 *HadronPhysics2* consortium







STARTING POINT  
OF  
DYNAMICAL MODELS




STARTING POINT  
OF  
STATISTICAL MODELS



fragments ( $Z > 2$ )  
liquid drops

At freeze-out : thermal and chemical equilibrium



Production of hypernuclei in relativistic heavy ion collisions


- production of many hyperons
- multiple coalescence of hyperons with fragments
- $(\pi, K)$ ,  $(K, \pi)$  and  $(K^-, K^+)$  reactions on fragments

# Concept of the HypHI experiment

Produced hypernucleus at similar velocity of projectiles



large Lorentz factor:  $\gamma > 3$ : life-time 200 ps  $\rightarrow$   $\sim 600$  ps

hypernuclear decay in flight




# Monte Carlo simulations

	Expected cross section [ $\mu\text{b}$ ]	Reconstructed events /week
${}^3_{\Lambda}\text{H}$	0.1	$2.8 \times 10^3$
${}^4_{\Lambda}\text{H}$	0.1	$2.6 \times 10^3$
${}^5_{\Lambda}\text{He}$	0.5	$6.5 \times 10^3$



# Scintillating fibre detectors

- **fibre diameter: 0.83 mm**
- **HAMAMATSU H7260KS**
- **X and Y tracking**
  - Position resolution:  $\sim 0.5$  mm
- **discriminator cards (1400 ch) from KAOS**
- **energy readout by CAEN QDC for TR0**
- **time readout by VUPROM 2**








anti-proton beam induced hypernuclei production:

- high resolution  $\gamma$ -spectroscopy of double  $\Lambda\Lambda$  hypernuclei
- weak decays

# General Idea

Use  $\bar{p}p$  Interaction to produce a hyperons which are tagged by the anti-hyperon or its decay products



# Double hypernuclei with kaons

**Indirect reaction  
(quasi-free in nucleus):**




$p(K^-) \approx 1.8$  (1.66) GeV/c  $\rightarrow$   
 $p(K^+, \Xi^-) \approx 1.3, 0.5$  (1.15, 0.5)  
 GeV/c (forward)

**Direct reactions:**




$\sigma(\theta=0) \approx 3.5 \mu\text{b/sr}$  (Dover & Gal)

$p(K^-) \approx 1.8$  (1.66) GeV/c  $\rightarrow p(K^+) \approx 1.39$  (1.24) GeV/c (forward)



$\sigma(\theta) \approx 10 \text{ nb/sr}$  (May)

$p(K^-) \approx 1.8$  (1.66) GeV/c  $\rightarrow p(K^+) \approx 1.42$  (1.28) GeV/c (forward)



# Double hypernuclei with antiprotons

## ***Indirect reaction:***



(PANDA)

## ***Two-step process in one nucleus:***



(FLAIR)

## ***Recoilless production:***






## Formation of double hypernuclei from cascade particles

1.  $dE(\Xi)/dx \rightarrow \text{stop} + \text{capture}$
2. *hyperatom + atomic decay*
3. *capture in nucleus ( ${}^{A'}_{\Xi}Z'$ )*
4. *conversion:  $\Xi + p \rightarrow \Lambda\Lambda + 28\text{MeV}$*
5. *double hypernucleus ( ${}^{A'}_{\Lambda\Lambda}Z'$ )*

# Production mechanism at FLAIR

[original idea K. Kilian 1987]



# GSI and FAIR facilities


## **GSI, Darmstadt**

- heavy ion physics
- nuclear structure
- atomic and plasma physics
- cancer therapy

**PANDA**

## **FAIR: New facility**

- heavy ion physics
- higher intensities & energies
- antiproton physics



**New**

Existing

## Primary Beams

- $^{238}\text{U}^{28+}$  :  $10^{12}/\text{s}$  @ 1.5-2 AGeV;
- $^{238}\text{U}^{92+}$  :  $10^{10}/\text{s}$  @ up to 35 AGeV
- **Protons** :  $2 \times 10^{13}/\text{s}$  @ 30 GeV; up to 90 GeV
- 100-1000 times present intensity

## Secondary Beams

- broad range of radioactive beams up to 1.5 - 2 AGeV
- intensity up to 10 000x over present
- antiprotons 0 - 15 GeV


## Storage and Cooler Rings

- radioactive beams
- $e^-$  - A (or Antiproton-A) collider
- $10^{11}$  stored and cooled antiprotons 0.8 - 14.5 GeV/c
- *In future*: polarized antiprotons (?)

## Key Technical Features

- cooled beams
- rapidly cycling superconducting magnets
- parallel operation

# High Energy Storage Ring



## H E S R Performance

Racetrack shaped ring: 574 m length

Luminosity/Intensity:

- Pbar production rate:  $2 \times 10^7$  /s
- High luminosity mode:  $L = 2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
- High resolution mode:  $L = 2 \times 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$   
(for target thickness  $4 \times 10^{15} \text{ atoms/cm}^2$ )



Momentum range:

- 1.5 – 15 GeV/c (0.831- 14.1 GeV)
- Revolution frequency:  $5 \times 10^5$  Hz

Momentum resolution:

- High luminosity mode:  $\Delta p/p = 10^{-4}$   
(stochastic cooling above 3.8 GeV/c)
- High resolution mode:  $\Delta p/p = 10^{-5}$   
(electron cooling)

# Beam cooling in HESR




## Resonance scan:

- Energy resolution down to  $\sim 50$  keV
- Tune  $E_{CM}$  to probe resonance
- Get precise mass and width

$E_{CM}$

# Production mechanism at PANDA







- $\bar{p}$  atoms: x-rays
- conversion:
 
$$\bar{p}(dss) p(uud) \rightarrow \Lambda(uds) \Lambda(uds)$$

$$\Delta Q = 28 \text{ MeV}$$
 conversion probability  $\sim 5\text{-}10\%$



[Y. Hirate et al., Nucl. Phys. **A639**, 389c (1998),  
 Y. Hirate et al., Prog. Theor. Phys. **102**, 89 (1999)]

# Detection and triggering



Performance of HPGe in strong B-field: EU FP6 HadronPhysics project



# Recoil momentum



# Application to target design

**First step:  
stop  $\Xi^-$  particles  
in nuclear targets**

**optimization  
of geometry**







**stopping points  
concentrated  
between  $40^\circ$  and  
 $90^\circ$  in a radial  
thickness of 20 mm**

# Preparatory experiment in 2004-5

experimental set-up:


- ALADiN dipole magnet
- Co source of 370 kBq
- VEGA clover and
- Euroball cluster




supported by a 6<sup>th</sup> European Framework  
HadronPhysics I3 Joint Research Activity

# Hypernuclear set-up at PANDA



- $\theta_{\text{lab}} < 45^\circ$ :  $\Xi$ -bar, K trigger and PID in PANDA spectrometer
- $\theta_{\text{lab}} = 45^\circ$ - $90^\circ$ :  $\Xi$ -capture and hypernuclei formation
- $\theta_{\text{lab}} > 90^\circ$ :  $\gamma$ -detection with HPGe at backward angles



# The PANDA PID detectors



# TOF for low momentum kaons



► Scintillating fibers (START)

~2000 fibers placed in two rings ⊕ readout with SiPM

► TOF barrel (STOP)

time resolution ~ 80 ps with 16 slabs

[simulations by A. Sanchez, U Mainz]

# Comparison of experiments

<i>experiment</i>	<i>reaction</i>	<i>device</i>	<i>beam/ target</i>	<i>status</i>
BNL-AGS E885	$(\Xi^-, ^{12}\text{C}) \rightarrow \begin{matrix} ^{12}\text{B} \\ \Lambda\Lambda \end{matrix} + n$	neutron detector arrays	K <sup>-</sup> beam, diamond target	20,000 <b>stopped</b> $\Xi^-$
BNL-AGS E906	$2\pi$ decays	cylindrical detector system	K <sup>-</sup> beam	few tens $2\pi$ decays of $^4_{\Lambda\Lambda}\text{H}$
KEK-PS E373	$(\text{K}^-, \text{K}^+)\Xi$	emulsion	$(\text{K}^-, \text{K}^+)$	several hundred stopped $\Xi^-$
<i>facility</i>	<i>reaction</i>	<i>device</i>	<i>beam/ target</i>	<i>captured</i> $\Xi^-$ <i>per day</i>
JHF	$(\text{K}^-, \text{K}^+)\Xi$	spectrometer, $\Delta\Omega = 30$ msr	$8 \cdot 10^6/\text{sec}$ 5 cm $^{12}\text{C}$	< 7,000 expected
cold anti- protons	$p \bar{p} \rightarrow \text{K}^* \bar{\text{K}}^*$ $\bar{\text{K}}^* \text{N} \rightarrow \Xi \text{K}$	vertex detector	$10^6$ stopped $\bar{p}$ per sec	2,000 expected
PANDA	$p \bar{p} \rightarrow \Xi \bar{\Xi}$	vertex detector + $\gamma$ -array	$L = 2 \cdot 10^{32}$ , thin target, production & decay target	<b>3,000</b> „golden events“ expected <b>~ 300,000</b> KK trigger expected

# Estimated rates

## “Golden events”:

luminosity  $2 \cdot 10^{32} \text{ cm}^{-2}\text{s}^{-1}$

$\Xi^+\Xi^-$  cross section 2mb for pp  $\Rightarrow$  700 Hz

p(100-500 MeV/c)  $p_{500} \approx 0.0005$

$\Xi^+$  reconstruction probability 0.5

stopping and capture probability  $p_{\text{CAP}} \approx 0.20$

total captured  $\Xi^- \Rightarrow$  3000 / day

$\Xi^-$  to  $\Lambda\Lambda$ -nucleus conversion probability  $p_{\Lambda\Lambda} \approx 0.05$

total  $\Lambda\Lambda$  hypernucleus production  $\Rightarrow$  4500 / month

gamma emission/event,  $p_{\gamma} \approx 0.5$

$\gamma$ -ray peak efficiency  $p_{\text{GE}} \approx 0.1$

~ 7/day „golden“  $\gamma$ -ray events with  $\Xi^+$  trigger


~ 700/day with *KK* trigger

# Summary

Hypernuclei offer a bridge between traditional nuclear and hadron physics

It helps to explore fundamental questions

- how do nucleons and nuclei form out of quarks?
- can nuclear structure be derived *quantitatively* from QCD?
- properties of strange baryons in nuclei and structure of QCD vacuum?



---

Thank you!